

LETTER TO THE EDITOR

Two very nearby ($d \sim 5$ pc) ultracool brown dwarfs detected by their large proper motions from WISE, 2MASS, and SDSS data^{*}

R.-D. Scholz, G. Bihain, O. Schnurr, and J. Storm

Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany
e-mail: rdscholz@aip.de, gbiain@aip.de, oschnurr@aip.de, jstorm@aip.de

Received May 20, 2011; accepted ...

ABSTRACT

Aims. WISE provides an infrared all-sky survey which aims at completing our knowledge on the possibly dramatically increasing number of brown dwarfs with lower temperatures. We search for the nearest representatives of the coolest brown dwarfs, which will be very interesting for detailed follow-up observations, once they have been discovered.

Methods. We have used the preliminary data release from WISE, selected bright candidates with colours typical of late-T dwarfs, tried to match them with faint 2MASS and SDSS objects, to determine their proper motions, and to follow-up them spectroscopically.

Results. We have identified two new ultracool brown dwarfs, WISE J0254+0223 and WISE J1741+2553, with large proper motions of about 2.5 and 1.5 arcsec/yr, respectively. With their $w_1-w_2 \sim 3.0$ and $J-w_2 \sim 4.0$ colour indices we expect them both to have a spectral type of $\sim T9-T10$ and absolute magnitude of $M_{w_2} \sim 14$. For WISE J1741+2553 we confirm a spectral type of T10 from near-infrared spectroscopy with LBT/LUCIFER1. From their bright WISE w_2 magnitudes of 12.7 and 12.3, we estimate distances of about 5.5 pc and 4.6 pc and tangential velocities of ~ 65 km/s and ~ 34 km/s indicating Galactic thick and thin disk membership, respectively.

Key words. Astrometry – Proper motions – Stars: distances – Stars: kinematics and dynamics – brown dwarfs – solar neighbourhood

1. Introduction

The Wide-field Infrared Survey Explorer (WISE; Wright et al. 2010) observed the sky in four infrared bands (w_1 at $3.4 \mu\text{m}$, w_2 at $4.6 \mu\text{m}$, w_3 at $12 \mu\text{m}$, and w_4 at $22 \mu\text{m}$). It allows the detection of nearby cool brown dwarfs (spectral types $>T5$) with much higher efficiency than the existing Two Micron All Sky Survey (2MASS; Skrutskie et al. 2006) and the ongoing Sloan Digital Sky Survey with its two recent data releases (SDSS DR7; Abazajian et al. 2009, and SDSS DR8; Aihara et al. 2011).

As for the nearest stars in the catalogue of Gliese & Jahreiß (1991) we expect the nearest cool brown dwarfs to be high proper motion (HPM) objects. Fig. 1 shows the correlation between proper motion and parallax. HPM stars with $\mu > 1$ arcsec/yr lie typically within 10 pc. Compared to 912 nearby stars shown with their mean values, only few (34) late-T dwarfs with parallax measurements are available. However, seven out of ten HPM late-T dwarfs with $\mu > 1$ arcsec/yr fall in the 10 pc sample, whereas the other three are very close to the 10 pc horizon.

This motivated our HPM search for cool brown dwarfs in the immediate solar neighbourhood using the preliminary WISE data release combined with the previous near-infrared (2MASS) and deep optical (SDSS) surveys. First results are presented.

2. Selection of candidates and cross-identification

To identify possible HPM and thus nearby cool brown dwarfs in the WISE data we used the following selection criteria:

- $w_1-w_2 > 2$ and $w_2-w_3 < 2.5$ - these are the colour constraints also applied by Burgasser et al. (2011) in search of late-T

^{*} based on observations with the Large Binocular Telescope (LBT)




Fig. 1. Mean distances of nearby stars (from Gliese & Jahreiß 1991) with 10% accurate parallaxes vs. their proper motions. The number of stars in each proper motion bin is given on the top. The last two bins are wider to have enough stars per bin. Error bars show the width of the distribution in each bin. Known $>T5$ dwarfs with parallaxes (from the compilation of Gelino et al. 2011, supplemented by data from Marocco et al. 2010, Liu et al. 2011, Luhman, Burgasser & Bochanski 2011, and Subasavage et al. 2009) are overplotted as red crosses.

dwarfs, where the second condition aims at excluding extragalactic sources (Wright et al. 2010).

- $w_2 < 13$ - we considered only bright WISE sources, since those have a higher probability to show up in 2MASS or SDSS if we consider the limiting magnitudes of these surveys (see Table 9 in Metchev et al. 2008) and the typi-

cal colours of known late-T dwarfs ($J-w2 \gtrsim 2$; Mainzer et al. 2011 and $z-J \sim 3.5$; Metchev et al. 2008)

- We included only WISE point sources outside the Galactic plane ($|b| > 10^\circ$ to avoid problems due to reddening.
- We looked only for WISE sources without a 2MASS counterpart within 3 arcsec, which implies a minimum proper motion of $\mu \gtrsim 0.3$ arcsec/yr with the typical epoch difference of 10 years between 2MASS and WISE.

Only 98 candidates remained after this selection, and all of them were subsequently inspected by eye on 2×2 arcmin finding charts (as shown in Fig. 2) from WISE, 2MASS, and SDSS (if available) in search of shifted counterparts with corresponding proper motions up to about 6 arcsec/yr. The majority of candidates were rejected as reddened stars, background galaxies, and artefacts (in diffraction spikes of bright stars), and for a few candidates no lucid explanation could be given. However, four known T7-T8 dwarfs (see Gelino et al. 2011 and references therein; with proper motions from 1.3 to 2 arcsec/yr), 2MASSI J0415195–093506 (T8 at 5.7 pc), 2MASSI J0727182+171001 (T7 at 9.1 pc), 2MASS J07290002–3954043 (T8p), and Gliese 570D (T7.5 at 5.9 pc) were found and two objects newly discovered. The two new objects are described in the next section.

3. Discovery of two new HPM brown dwarfs

The four known late-T dwarfs, which we identified on the WISE and 2MASS finding charts with positional separations of about 15 to 25 arcsec, showed up as bright and unusual ‘green’ sources in the WISE false colour composites and as relatively faint ($15.3 < J < 15.9$) and ‘blue’ ($-0.2 < J-H < +0.2$ and $0.0 < J-K_s < +0.3$) 2MASS sources. Except for the T8p dwarf, they were still detected in the 2MASS K_s -band. For two new ‘green’ and ‘yellow’ WISE sources (see Fig. 2), WISEPC J025409.45+022359.1 (hereafter WISE J0254+0223) and WISEPC J174124.25+255319.5 (hereafter WISE J1741+2553), we found even fainter ($16.4 < J < 16.6$) 2MASS counterparts at similar large separations. The latter two objects are not detected in the 2MASS K_s -band, and only the second one is blue ($J-H \approx +0.1$ and $J-K_s \lesssim -0.3$), whereas the first one appears to be relatively red ($J-H \approx +0.7$ and $J-K_s \lesssim +0.6$). However, the 2MASS magnitudes of the two new objects are close to the survey limits, and the corresponding colours from 2MASS are rather uncertain, compared to those of the four brighter known late-T dwarfs.

The additional identification of WISE J0254+0223 in SDSS DR8 (Aihara et al. 2011) and of WISE J1741+2553 in SDSS DR7 (Abazajian et al. 2009) led to clear proper motion fits (Fig. 3). The very large proper motions are a first hint that these objects should be very close to the Sun. Both objects are only detected in the SDSS z -band which is typical of nearby late-T dwarfs. Metchev et al. (2008) mentioned that there were no T8 and later type brown dwarfs known in the SDSS. The 11 late-T dwarf candidates identified by Scholz (2010) in SDSS and UKIDSS (six of which have been confirmed spectroscopically as T5-T8 dwarfs by Burningham et al. 2010, Burgasser et al. 2010, and Murray et al. 2011) have $19.9 < z < 20.7$. None of them were detected in 2MASS. In comparison, our two new objects are with $z=19.9$ and $z=19.7$ relatively bright and have at least 2-3 times larger proper motions than the above mentioned 11 objects.

The photometry and different epoch positions of WISE J0254+0223 and WISE J1741+2553 from SDSS, 2MASS and WISE, as well as their proper motions are shown in




Fig. 2. Finding charts of WISE J0254+0223 (left column) and WISE J1741+2553 (right column) showing 2×2 arcmin fields centered on the WISE positions with North up and East to the left. The charts are sorted by the epochs of the images, 2MASS, SDSS, WISE (from top to bottom). The lower row shows WISE false colour composites.

Table 1. Note that the proper motions are derived from simple linear fitting and may be affected by parallactic motions.


4. Photometric classification and distances

With their large $w1-w2$ colour indices of +3.04 and +2.92 WISE J0254+0223 and WISE J1741+2553 are expected to have spectral types of $\sim T7$ - $T10$ according to Mainzer et al. (2011), although we find for the four known T7-T8 dwarfs detected in our HPM search (Sect. 1) somewhat smaller values of $+2.30 < w1-w2 < +2.85$. Therefore, we believe our two new objects are more likely ultracool ($\gtrsim T8$) brown dwarfs. In support of this classification we mention their large $J-w2$ indices of +3.85 and +4.14 (compared to values from +2.67 to +3.46 for the four known T7-T8 dwarfs) corresponding to an abso-


Table 1. Astrometry & photometry of two new HPM brown dwarfs

Parameter	WISE J0254+0223	WISE J1741+2553
WISE RA (J2000)	02 54 09.4501	17 41 24.2578
WISE Dec (J2000)	+02 23 59.136	+25 53 19.507
WISE epoch	2010.182	2010.182
SDSS RA (J2000)	02 54 09.2449	17 41 24.4306
SDSS Dec (J2000)	+02 23 58.268	+25 53 27.805
SDSS epoch	2008.764	2004.7095
SDSS run/data release	7717/DR8	4832/DR7
2MASS RA (J2000)	02 54 07.8864	17 41 24.6206
2MASS Dec (J2000)	+02 23 56.346	+25 53 34.361
2MASS epoch	2000.723	2000.277
$\mu_\alpha \cos \delta$ [mas/yr]	+2496 \pm 46	-492 \pm 43
μ_δ [mas/yr]	+276 \pm 47	-1500 \pm 11
SDSS z [mag]	19.861 \pm 0.074	19.745 \pm 0.105
2MASS J [mag]	16.557 \pm 0.156	16.451 \pm 0.099
2MASS H [mag]	15.884 \pm 0.199	16.356 \pm 0.216
2MASS K_s [mag]	>16.006	>16.785
WISE $w1$ [mag]	15.743 \pm 0.070	15.228 \pm 0.040
WISE $w2$ [mag]	12.707 \pm 0.031	12.312 \pm 0.025
WISE $w3$ [mag]	11.042 \pm 0.131	10.675 \pm 0.075

Notes. We have also detected both objects at the positions 02 54 07.301 +02 23 56.91 (1996.805) and 17 41 24.649 +25 53 40.25 (1996.449), respectively on I -band photographic Schmidt plates (not shown in Fig. 2) scanned in the SuperCOSMOS Sky Surveys (Hambly et al. 2001). The proper motions involving these positions (+2438.7 \pm 40.6, +176.1 \pm 64.1 and -407.8 \pm 59.7, -1507.2 \pm 7.0, respectively) confirm our HPM measurements but are slightly less accurate than our preferred solutions given in the table above and shown in Fig. 3.


Fig. 3. Relative positions from 2MASS, SDSS, and WISE (from left to right) and proper motion fits for WISE J0254+0223 (filled lozenges, solid lines) and WISE J1741+2553 (filled squares, dashed lines)

lute magnitude of $M_{w2} \sim 14$ if we use the linear relation between M_{w2} and $J-w2$ given in Wright et al. (2011). For comparison, UGPS J072227.51-054031.2 (hereafter UGPS J0722-0540), the T10 dwarf at 4.1 pc (Lucas et al. 2010), is detected in WISE with $w2=12.17$ and has an absolute magnitude of $M_{w2}=14.11$, whereas all T dwarfs with known distances from Patten et al. (2006) plotted by Wright et al. (2011) have brighter absolute magnitudes ($M_{w2} < 13.5$) with colour indices $J-w2 < +3.3$. The $J-w2$ colours of WISE J0254+0223 and WISE J1741+2553 do not reach the very large values (≥ 5) of the two coolest known (probably >T10) but more distant


Fig. 4. LBT/LUCIFER1 $H+K$ - (bottom) and J -band (top) spectra of WISE J1741+2553 and comparison objects (see Table 2).

brown dwarfs WD 0806-661B at ~ 19 pc (Luhman, Burgasser & Bochanski 2011; Rodriguez et al. 2011; Subasavage et al. 2009) and CFBDSIR J145829+101343B at ~ 23 pc (Liu et al. 2011) but are comparable with that of the A component (T9.5) of the latter as shown in Wright et al. (2011).

For a first approximation of the distances of WISE J0254+0223 and WISE J1741+2553, we use the preliminary classification of both objects as $\sim T9$ -T10 dwarfs with an absolute magnitude of $M_{w2} \sim 14$. We may, however, assume a large uncertainty of 0.5 mag in this absolute magnitude, since for the brighter T dwarfs shown in Wright et al. (2011) there is a similar large spread in absolute magnitudes. We then arrive at estimated distances from the Sun for WISE J0254+0223 and WISE J1741+2553 of $5.5^{+1.4}_{-1.1}$ pc and $4.6^{+1.2}_{-1.0}$ pc, respectively.

With these distance estimates, we compute absolute magnitudes $M_z \sim 21.2$ and $M_z \sim 21.4$, respectively for WISE J0254+0223 and WISE J1741+2553. These values are 1.3-1.5 mag fainter than the $M_z \sim 19.9$ of Ross 485C (=Hip 63510C), the only T8 dwarf with distance and SDSS measurements now available (Goldman et al. 2010; Scholz 2010; Burgasser et al. 2010). The latter is in good agreement with the assumed $M_z \sim 20.0$ for T8 dwarfs in Metchev et al. (2008). On the other hand, the z -band photometry obtained by Lucas et al. (2010) for the T10 dwarf UGPS J0722-0540 leads to M_z values between 22.4 and 22.6, about 1 mag fainter than our values. The intermediate M_z values of our new objects support their classification as $\sim T9$ -T10 dwarfs.

5. Spectroscopic observations with LBT/LUCIFER

We have used the Large Binocular Telescope (LBT) near-infrared spectrograph LUCIFER1 (Mandel et al. 2008; Seifert et al. 2010; Ageorges et al. 2010) in long slit spectroscopic mode with the $H+K$ (200 lines/mm + order separation filter) and $zJHK$ gratings (210 lines/mm + J filter) to observe

Table 2. Spectral indices obtained from LBT/LUCIFER1 spectra (SpT from comparison with T9 average from Lucas et al. 2010 in brackets)

Object	SpT	Reference for SpT	W_J	$\text{NH}_3 - \text{H}$	$\text{H}_2\text{O} - \text{H}$	$\text{CH}_4 - \text{H}$	$\text{CH}_4 - \text{K}$
Ross 485C	T8	Burgasser et al. 2010	0.264 (<T9)	0.677 (<T9)	0.201 (<T9)	0.109 (<T9)	0.055 (>T9)
UGPS J0722–0540	T10	Lucas et al. 2010		0.520 (>T9)	0.138 (<T9)	0.075 (>T9)	0.079 (>T9)
WISE J1741+2553	T10	this paper	0.234 (>T9)	0.551 (<T9)	0.119 (>T9)	0.066 (>T9)	0.056 (>T9)

UGPS J0722–0540 on 2011 March 06 (only in $H+K$, total integration 30 min) and WISE J1741+2553 (40 min $H+K$, 27 min J) together with Ross 485C (30 min $H+K$, 20 min J) on 2011 May 12. The central wavelengths were chosen at $1.835 \mu\text{m}$ ($H+K$) and $1.25 \mu\text{m}$ (J) yielding a coverage of $1.38\text{--}2.26$ and $1.17\text{--}1.32 \mu\text{m}$, respectively. The slit width was of 2 arcsec for UGPS J0722–0540 and 1 arcsec (4 pixels) for the other objects. For the 1 arcsec slit, the spectral resolving power is $R=\lambda/\Delta\lambda \approx 4230, 940$, and 1290 at $\lambda \approx 1.24, 1.65$, and $2.2 \mu\text{m}$, respectively. Observations consisted of individual 75 s exposures in $H+K$ and 200 s exposures in J with shifting the target along the slit following an ABBA pattern (to allow sky subtraction) until the total integration time was reached. A0V standards were observed just before or after the targets with similar airmass.

The raw spectroscopic data were reduced using standard routines within the IRAF¹ environment. The spectra were sky-subtracted, aligned, combined, optimally extracted, and wavelength-calibrated using vacuum wavelengths of Ar arc lamps and the deep telluric absorption features. No flat division was applied to avoid including additional noise features from the halogen lamp. The instrumental response and telluric bands were removed by dividing by the A0 star spectra and multiplying by a black-body spectrum with the same effective temperature. The intrinsic lines in the hot-star spectra were removed before dividing the science spectra. Residuals from intense sky emission lines in the target spectra were removed by interpolation across the lines. The J and $H+K$ spectra were normalized by the average flux in the range $1.20\text{--}1.30$ and $1.52\text{--}1.61 \mu\text{m}$, respectively.

The $H+K$ spectrum of WISE J1741+2553 is very similar to that of the T10 dwarf UGPS J0722–0540, whereas both spectra ($H+K$ and J) of WISE J1741+2553 indicate a later type if compared to those of the T8 dwarf Ross 485C (Fig. 4). We have computed spectral indices used for the classification of ultracool brown dwarfs (see Burningham et al. 2008 and references therein) and compare them with average values of T9 dwarfs given by Lucas et al. (2010) in Table 2. Unfortunately, no other indices than W_J could be determined in the J -band, since the LUCIFER J grating provides a very narrow wavelength interval. Our indices are in good agreement with the measurements of Lucas et al. (2010) for UGPS J0722–0540 and of Burgasser et al. 2010 for Ross 485C, except for $\text{NH}_3 - \text{H}$, where our indices are larger by about 0.03 and 0.04, respectively. As in the case of the known T10 dwarf, all but one indices of WISE J1741+2553 classify it as >T9. Therefore we assign a spectral type of T10.

6. Conclusions

1. While WISE J0254+0223 and WISE J1741+2553 are likely similar to the few other T9–T10 brown dwarfs known, they are the first ultracool brown dwarfs detected in both 2MASS and SDSS. With their relatively bright magnitudes they are

excellent targets for detailed spectroscopic investigations and for high resolution imaging in search of possible binarity. They may become important benchmark sources at the boundary between the T-type and the suggested Y-type (Kirkpatrick et al. 1999) classes of brown dwarfs.

2. The photometric classification of both objects as \sim T9–T10 dwarfs was confirmed for WISE J1741+2553, for which we obtained spectroscopy with LBT/LUCIFER1 and classified it as an ultracool T10 dwarf. This is in all likelihood true for WISE J0254+0223, too.
3. Our photometric distance estimates place both objects at about 5 pc from the Sun, probably nearly as close as the T10 dwarf UGPS J0722–0540 in the Galactic plane ($b=+4^\circ$), and make them highly interesting targets for near-infrared parallax programs. They fall still clearly in the 10 pc sample if we use the upper limits of our distance estimates.
4. The larger proper motion and slightly larger distance of the higher Galactic latitude object WISE J0254+0223 ($b=-48^\circ$) lead to a relatively high tangential velocity of ~ 65 km/s (possibly indicating a Galactic thick disk membership), compared to ~ 34 km/s for WISE J1741+2553 ($b=+26^\circ$), which is typical of the thin disk population.

Acknowledgements. The authors would like to thank Roland Gredel, Jochen Heidt, Jaron Kurk, Ric Davies, and all observers at the LBT for their assistance and help during the preparation and execution of the LUCIFER observations.

This research has made use of the NASA/IPAC Infrared Science Archive, which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration, and of data products from WISE, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration, from 2MASS, and from SDSS DR7 and DR8. Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy. The SDSS-III web site is <http://www.sdss3.org/>.

References

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, *ApJS*, 182, 54
- Ageorges, N., Seifert, W., Jütte, M., et al. 2010, *SPIE*, 7735, 53
- Aihara, H., Allende Prieto, C., An, D., et al. 2011, *ApJS*, 193, 29
- Burningham, B., Pinfield, D. J., Leggett, S. K., et al. 2008, *MNRAS*, 391, 320
- Burningham, B., Pinfield, D. J., Lucas, P. W., et al. 2010, *MNRAS*, 406, 1885
- Burgasser, A. J., Simcoe, R. A., Bochanski, J. J., et al. 2010, *ApJ*, 725, 1405
- Burgasser, A. J., Cushing, M. C., Kirkpatrick, J. D., et al. 2011, *arXiv:1104.2537*
- Gelino, C. R., Kirkpatrick, J. D., & Burgasser, A. J. 2011, online database for 804 L and T dwarfs at DwarfArchives.org (status: 14 February, 2011)
- Gliese, W., & Jahreiß, H. 1991, On: The Astronomical Data Center CD-ROM: Selected Astronomical Catalogs, Vol. I; L.E. Brothmann, S.E. Gesser (eds.), NASA/Astronomical Data Center, Goddard Space Flight Center
- Goldman, B., Marsat, S., Henning, T., Clemens, C., & Greiner, J. 2010, *MNRAS*, 405, 1140
- Hambly, N. C., MacGillivray, H. T., Read M. A., et al. 2001, *MNRAS*, 326, 1279
- Kirkpatrick, J. D., Reid, I. N., Liebert, J., et al. 1999, *ApJ*, 519, 802
- Liu, M. C., Delorme, P., Dupu, T. J., et al. 2011, *arXiv:1103.0014*
- Lucas, P. W., Tinney, C. G., Burningham, B., et al. 2010, *MNRAS*, 408, L56
- Luhman, K. L., Burgasser, A. J., & Bochanski, J. J. 2011, *ApJ*, 730, L9
- Mandel, H., Seifert, W., Hofmann, R., et al. 2008, *SPIE*, 7014, 124
- Mainzer, A., Cushing, M. C., Skrutskie, M. F., et al. 2011, *ApJ*, 726, 30
- Marocco, F., Smart, R. L., Jones, H. R. A., et al. 2010, *A&A*, 524, A38

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

- Metchev, S. A., Kirkpatrick, J. D., Berriman, G. B., & Looper, D. 2008, *ApJ*, 676, 1281
- Murray, D. N., Burningham, B., Jones, H. R. A., et al. 2011, arXiv:1101.4881
- Patten, B. M., Stauffer, J. R., Burrows, A., et al. 2006, *ApJ*, 651, 502
- Rodriguez, D. R., Zuckerman, B., Melis, C., & Song, I. 2011, *ApJ*, 732, L29
- Scholz, R.-D. 2010, *A&A*, 515, A92
- Seifert, W., Ageorges, N., Lehmitz, M., et al. 2010, *SPIE*, 7735, 256
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Subasavage, J. P., Jao, W.-Ch., Henry, T. J., et al. 2009, *AJ*, 137, 4547
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868
- Wright, E. L., Mainzer, A., Gelino, C., & Kirkpatrick, D. 2011, arXiv:1104.2569